arXiv:1403.4741v1 [math.CO] 19 Mar 2014

DIAMETER 2 CAYLEY GRAPHS OF DIHEDRAL GROUPS

GRAHAME ERSKINE

Abstract. We consider the degree-diameter problem for Cayley graphs of dihedral groups. We find upper and lower bounds on the maximum number of vertices of such a graph with diameter 2 and degree d. We completely determine the asymptotic behaviour of this class of graphs by showing that both limits are asymptotically $d^2/2$.

1. Introduction

The *degree-diameter problem* seeks to determine the largest possible number $n(d,k)$ of vertices of a graph of maximum degree d and diameter k. The survey [3] summarises current known results for the general case and also for various restricted problems where we consider only graphs of certain classes. In this paper we restrict attention to Cayley graphs of dihedral groups with diameter 2 and prove an asymptotic limit for the maximum order of such graphs.

An elementary counting argument yields an upper limit of d^2+1 (the well-known *Moore bound*) for a graph of maximum degree d and diameter 2. Thus for any family of diameter 2 graphs the largest possible asymptotic order is d^2. When we restrict consideration to Cayley graphs, it is known [4] that there is a relatively sparse family of groups (affine groups over finite fields of characteristic 2) on which Cayley graphs can be constructed with asymptotic order d^2. For more elementary families of groups, less is known. For example an extension of the counting argument shows that for abelian groups the largest possible asymptotic order is $d^2/2$, but the best known result [2] has limit $3d^2/8$. We show that the asymptotic limit for dihedral groups is precisely $d^2/2$, first by obtaining a lower bound by way of a construction involving Galois fields, and then by finding an upper bound for generalised dihedral groups by a counting argument.

We denote the dihedral group of order n by D_n. We will view the usual dihedral group as an example of a *generalised dihedral group* $G \rtimes C_2$ which is a semidirect product of an abelian group G with the multiplicative group $\{\pm 1\}$ where the action on G is via its inversion automorphism. For a group G and a subset $S \subset G$ which is inverse-closed and does not contain the identity in G, we define the Cayley graph $\mathrm{Cay}(G,S)$ to have vertex set G and an undirected edge gh if and only if $gh^{-1} \in S$. The graph is regular of

2010 *Mathematics Subject Classification.* 05C25.

Key words and phrases. Degree-diameter problem, Cayley graph, Dihedral group.

degree $d = |S|$. From the definition of Cayley graphs it is immediate that $\mathrm{Cay}(G, S)$ has diameter at most k if and only if each element of G can be expressed as a product of no more than k elements of the generating set S.

By $DC(d, k)$ we mean the largest number of vertices in a Cayley graph of a dihedral group having degree d and diameter k.

2. Results

Our first result uses a construction based on finite fields to obtain a lower bound for $DC(d, 2)$ for certain values of d. The method is similar to constructions in [2]. We also use the well-known result that the cyclic group $\mathbb{Z}_n$ has a diameter 2 Cayley graph with a generating set of size at most $2\lceil\sqrt{n}\rceil$. (To see see this, let $K = \lceil\sqrt{n}\rceil, M = \lfloor\frac{K}{2}\rfloor$ and take a generating set consisting of $\{\pm 1, \pm 2, \ldots, \pm M, \pm K, \pm 2K, \ldots, \pm MK\}$.)

Lemma 1. *If p is any prime and $d = 2(p + \lceil\sqrt{p}\rceil - 1)$, then $DC(d, 2) \geq 2p(p-1)$.*

Proof. Let F be the Galois field $GF(p)$ where p is a prime. The additive and multiplicative groups F^+ and F^* are cyclic of coprime orders so that $F^+ \times F^*$ is a cyclic group of order $n = p(p-1)$. Consider the dihedral group D_{2n} as a semidirect product $G = (F^+ \times F^*) \rtimes C_2$ where the cyclic group C_2 is thought of as the multiplicative group $\{\pm 1\}$ and acts on $F^+ \times F^*$ by inversion. Specifically, the multiplication rule is:

$$(a, b, c)(\alpha, \beta, \gamma) = (a + \alpha c, b\beta^c, c\gamma)$$

The subgroup $C = \langle(1, 1, 1)\rangle$ is cyclic of order p and so itself has a diameter 2 Cayley graph with respect to some generating set $\{c_i\}$ of cardinality $2\lceil\sqrt{p}\rceil$. Consider now a generating set S of the full group G containing:

$$\begin{aligned} v = (0, 1, -1) &\qquad (1 \text{ element}) \\ a_x = (0, x, 1), x \in F^* \setminus \{1\} &\qquad (p - 2 \text{ elements}) \\ b_x = (x, x, -1), x \in F^* &\qquad (p - 1 \text{ elements}) \\ c_i, i = 1 \ldots 2\lceil\sqrt{p}\rceil &\qquad (2\lceil\sqrt{p}\rceil \text{ elements}) \end{aligned}$$

It suffices to show that the set S is inverse-closed and that every element of the group can be expressed as the product of at most two of these generators.

Since $v^{-1} = v, a_x^{-1} = a_{x^{-1}}, b_x^{-1} = b_x$ and $\{c_i\}$ is inverse-closed it follows that S is inverse-closed. To show the diameter is 2 we consider all the possible cases as follows.

If $x \neq 0, x \neq y$ then $(x, y, -1) = (0, z, 1)(x, x, -1) = a_z b_x$ where $z = yx^{-1}$.

If $x \neq 0, x = y$ then $(x, y, -1) = (x, x, -1) = b_x$.

If $x = 0, y \neq 1$ then $(x, y, -1) = (0, y, 1)(0, 1, -1) = a_y v$.

If $x = 0, y = 1$ then $(x, y, -1) = (0, 1, -1) = v$.

If $y \neq 1, x \neq 0$ then $(x, y, 1) = (z, z, -1)(t, t, -1) = b_z b_t$ where $z = yx(y-1)^{-1}, t = x(y-1)^{-1}$.

If $y \neq 1, x = 0$ then $(x, y, 1) = (0, y, 1) = a_y$.

If $y = 1$ then $(x, y, 1) \in C$ and so is the product of at most two c_i. Since $|S| = 2(p + \lceil\sqrt{p}\rceil - 1)$ the result follows. □

The previous result shows that $\limsup_{d\to\infty} \frac{DC(d,2)}{d^2} \geq \frac{1}{2}$. The next result shows that 1/2 is in fact also an upper bound.

Lemma 2. *For any diameter 2 Cayley graph on a generalised dihedral group of order $2n$, the degree d satisfies $d \geq 2\sqrt{n} - 1$.*

Proof. Let $G = H \rtimes C_2$ where H is an abelian group of order n and C_2 acts on H by inversion. Suppose S is an inverse-closed subset of G such that $\mathrm{Cay}(G, S)$ has diameter 2. Let C be the index 2 subgroup of G isomorphic to H and write $S = A \cup B$ where $A \subset C$ and $B \subset G \backslash C$. Let $m_1 = |A|, m_2 = |B|$.

Consider how the n elements of $G \setminus C$ can be expressed as a product of at most two elements in S. There are m_2 possibilities from the set B itself, then $m_1 m_2$ elements of the form ab where $a \in A, b \in B$. Since $a^{-1}b = ba$ and the set A is inverse-closed the products of the form ba do not contribute any further elements. So we require:

$$m_2(m_1 + 1) \geq n$$

The degree d of the Cayley graph is $|S| = m_1 + m_2$. All numbers are inherently positive and so elementary calculus shows that the minimum possible value of $m_1 + m_2$ occurs when $m_2 = m_1 + 1 = \sqrt{n}$. So $d \geq 2\sqrt{n} - 1$. □

The bound in Lemma 2 is valid for all values of d, but as it stands Lemma 1 only holds for a restricted set of values. We can extend the result of Lemma 1 by using the ideas in [5] to obtain a lower bound valid for all values of d.

Lemma 3. *For all d, $DC(d, 2) \geq 2p(p - 1)$ where p is the largest prime satisfying $D(p) = 2(p + \lceil\sqrt{p}\rceil - 1) \leq d$.*

Proof. Let p be as in the statement, $n = 2p(p-1)$ and $G = D_n$. By Lemma 1 there is an inverse-closed subset $S \subset G$ with $|S| = D(p)$ such that $\mathrm{Cay}(G, S)$ has diameter 2. We can add $d - D(p)$ involutions from $G \setminus S$ to form a new inverse-closed generating set S'. The diameter of $\mathrm{Cay}(G, S')$ is still 2 and the result follows. □

Using the method of [5] we may use ideas from number theory to obtain a result independent of p. Specifically, from the result in [1] it follows that for all sufficiently large D, there is some prime p in the range $D - D^{0.525} \leq p \leq D$.

Theorem 4. *For all sufficiently large d, $DC(d, 2) \geq 0.5d^2 - 1.39d^{1.525}$.*

Proof. For given d, let p be the largest prime such that $2(p + \lceil\sqrt{p}\rceil - 1) \leq d$. Then p is at least as large as the largest prime q satisfying $2(q + \sqrt{q}) \leq d$. Rearranging this we find that q is the largest prime not exceeding $D =$

$\frac{1}{2}(d - \sqrt{2d+1} + 1)$. By the result of [1] for sufficiently large d we have $q \geq D - D^{0.525}$. So for large d we have:

$$\begin{aligned} p \geq q &\geq D - \left(\frac{d}{2}\right)^{0.525} \\ &= \frac{1}{2}\left(d - (2d+1)^{0.5} + 1 - 2^{0.475} d^{0.525}\right) \end{aligned}$$

For large d the term in $d^{0.525}$ dominates terms of lower powers of d and since $2^{0.475} \approx 1.389918$, for large d we have $p \geq \frac{1}{2}\left(d - 1.38992 d^{0.525}\right)$. For sufficiently large d we therefore have

$$2p(p-1) \geq \frac{d^2}{2} - 1.39 d^{1.525}$$

Since we can construct a Cayley graph of degree d and diameter 2 on the dihedral group of order $2p(p-1)$ by Lemma 3 the result follows. □

Lemma 2 and Theorem 4 allow us to determine completely the asymptotic behaviour of $DC(d, 2)$.

Corollary 5.

$$\lim_{d \to \infty} \frac{DC(d,2)}{d^2} = \frac{1}{2}$$

Remark. It is tempting to try to extend these results to other split extensions of abelian groups where the action is via an automorphism other than the inversion map. However the counting argument of Lemma 2 relies on the fact that in a Cayley graph $\mathrm{Cay}(G, S)$ the generating set S is closed under the inversion automorphism. The upper asymptotic bound $d^2/2$ would not necessarily hold for other more general semidirect products of an abelian group with C_2, although no family of such groups with a larger bound is known.

References

1. Roger C Baker, Glyn Harman, and János Pintz, *The difference between consecutive primes, II*, Proceedings of the London Mathematical Society **83** (2001), no. 3, 532–562.
2. Heather Macbeth, Jana Šiagiová, and Jozef Širáň, *Cayley graphs of given degree and diameter for cyclic, Abelian, and metacyclic groups*, Discrete Mathematics **312** (2012), no. 1, 94–99.
3. Mirka Miller and Jozef Širáň, *Moore graphs and beyond: A survey of the degree/diameter problem*, Electronic Journal of Combinatorics **DS14v2** (2013), 92pp.
4. Jana Šiagiová and Jozef Širáň, *Approaching the Moore bound for diameter two by Cayley graphs*, Journal of Combinatorial Theory, Series B **102** (2012), no. 2, 470–473.
5. Jozef Širáň, Jana Šiagiová, and Mária Ždímalová, *Large graphs of diameter two and given degree*, International Workshop on Optimal Network Topologies, Iniciativa Digital Politècnica, 2011, pp. 347–359.

Department of Mathematics and Statistics, The Open University, Milton Keynes, UK

E-mail address: Grahame.Erskine@open.ac.uk